




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C11D 1/65, 3/30, 1/86	A1	(11) International Publication Number: WO 96/12000 (43) International Publication Date: 25 April 1996 (25.04.96)
(21) International Application Number: PCT/US94/11779 (22) International Filing Date: 13 October 1994 (13.10.94) (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MORELLI, Joseph, Paul [US/US]; 541 Howell Street, Cincinnati, OH 45220 (US). LAPPAS, Dimitris [GR/BE]; 5, rue Alphons-Wauters, B-1020 Brussels (BE). RANDALL, Sherri, Lynn [US/US]; 11 Citadel Drive, Fairfield, OH 45014 (US). PANANDIKER, Rajan, Keshav [IN/US]; 6484 Oregon Pass, West Chester, OH 45069 (US). BOUTIQUE, Jean-Pol [BE/BE]; 32, rue Emile-Labarre, B-5030 Gembloux (BE). HOUSMEKERIDES, Chris, Efstathios [GR/BE]; 182, avenue Pois-de-Senteur, Boîte postale 3, B-1120 Brussels (BE). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i>

(54) Title: DETERGENT COMPOSITIONS CONTAINING AMINES AND ANIONIC SURFACTANTS

(57) Abstract

A liquid detergent composition comprising anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates, characterized in that said detergent composition further comprises an amine selected from i) primary amines having the formula: $R_1R_2R_3N$, wherein R_1 and R_2 are both H; and wherein R_3 is C_4-C_{18} , or $R_4X(CH_2)_n$, whereby X is -O-, -C(O)NH- or -NH-, R_4 is a C_6-C_{18} , n is between 1 to 5; ii) tertiary amines having the formula $R_1R_2R_3N$, wherein neither R_1 nor R_2 is H, R^1 and R^2 are C_1-C_8 alkylchains or (II), whereby n is between 1 to 5 and x is between 1 to 5, R_3 is either C_4-C_{18} or $R_4X(CH_2)_n$, whereby X is -O-, -C(O)NH- or -NH-, R_4 is a C_6-C_{18} , n is between 1 to 5 and R_5 is H or C_1-C_2 alkyl; and iii) mixtures of said primary and tertiary amines.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DETERGENT COMPOSITIONS CONTAINING AMINES
AND ANIONIC SURFACTANTS

Field of the invention

The present invention relates to detergent compositions containing surfactants selected from amines. More particularly, the invention is directed to detergent compositions containing anionic surfactants selected from the group consisting of alkyl alkoxyated sulfates and alkyl sulfates, said composition further containing specific primary and/or tertiary amines.

Background of the invention

Detergent compositions useful for cleaning purposes, such as laundering of fabrics, have commonly utilized a variety of surfactants.

The ability of surfactants to clean a large variety of soils and stains from fabrics present in the typical load of laundry is of high importance in the evaluation of detergent performance. Unfortunately, the relative ability of each surfactant to meet various performance criteria is among others depending on the presence of cosurfactants.

The recent trend towards partial or total replacement of Linear alkyl-benzene sulfonate surfactants (LAS) has urged the detergent formulators to rebalance their formulations with different surfactants. For example, quaternary ammonium salts are less efficient in boosting the greasy cleaning performance in Nil-LAS formulations.

There is thus a standing desire for performance and flexibility reasons to make available a surfactant system capable of providing optimum detergency performance which is equivalent to that of LAS-containing detergents.

The above objective has been met by a surfactant system comprising anionic surfactants selected from the group consisting of alkyl alkoxylated sulfates and alkyl sulfates, said surfactant system further comprising a cosurfactant selected from the group of primary or tertiary amines.

It has been surprisingly found that detergent compositions containing said surfactant system exhibit detergency performance equivalent to that of LAS-containing detergents.

In addition, it has been found that a small amount of certain primary and tertiary amines according to the present invention constitutes an efficient and compatible suds suppressing system. This additional benefit allows to use the amines in a dual function, e.g. surfactant and suds suppressor, thereby facilitating the formulation of concentrated liquid detergents.

Liquid detergent compositions formulated with said surfactant system are extremely useful when the liquid detergent compositions are in direct contact with the fabrics such as during pretreatment.

Amines have been described in the art in liquid detergent compositions. EP 160 762, EP 137 615 and EP 137 616 disclose liquid detergents which comprise cyclohexylamine. EP 177 165 discloses detergent compositions which comprise anionics, cellulase and a variety of primary, secondary, tertiary and quaternary amines. EP 11 340 discloses softener through the wash detergent compositions which comprise tertiary amines and clay. DE 32 07 487, GB 2 094 826, GB

2 095 275 and EP 137 397 disclose compositions which comprise anionics and quaternary amines. EP 120 528 discloses compositions comprising anionics, as well as tertiary amines. EP 26 528 and EP 26 529 disclose compositions comprising anionics and quaternary amines. Detergent compositions containing 1) primary or tertiary amines of the type useful herein in combination with 2) oleoyl sarcosinate are disclosed in U.S. Patent Application Serial No. 08/252,127, filed June 1, 1994. Compositions of this invention may or may not contain oleoyl sarcosinate surfactants.

Summary of the invention

The detergent compositions according to the present invention comprise anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates, characterized in that said detergent composition further comprises specific primary and/or a tertiary amines.

The detergent compositions preferably comprise at least 5%, more preferably from 10% to 65% and most preferably from 15% to 40% by weight of the surfactant system as described hereinabove.

Preferably, the weight ratio of the amine cosurfactant to the alkyl alkoxylated sulfates and alkyl sulfates is from 1:1 to 1:250, more preferably from 1:5 to 1:100 and most preferably from 1:10 to 1:40.

Detailed description of the Invention

The detergent compositions according to the present invention comprise anionic surfactants selected from the group of alkylalkoxy sulfates and alkyl sulfates.

Alkyl alkoxyated sulfates and/or alkyl sulfates

The alkyl alkoxyated sulfate surfactants hereof are water soluble salts or acids of the formula $RO(A)_mSO_3M$ wherein R is an unsubstituted C_{10} - C_{24} alkyl or hydroxyalkyl group having a C_{10} - C_{24} alkyl component, preferably a C_{12} - C_{18} alkyl or hydroxyalkyl, more preferably C_{12} - C_{15} alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are contemplated herein. Specific examples of substituted ammonium cations include ethanol-, triethanol-, methyl-, dimethyl-, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperidinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C_{12} - C_{15} alkyl polyethoxylate (1.0) sulfate (C_{12} - $C_{15}E(1.0)M$), C_{12} - C_{15} alkyl polyethoxylate (2.25) sulfate (C_{12} - $C_{15}E(2.25)M$), C_{12} - C_{15} alkyl polyethoxylate (3.0) sulfate (C_{12} - $C_{15}E(3.0)M$), and C_{12} - C_{15} alkyl polyethoxylate (4.0) sulfate (C_{12} - $C_{15}E(4.0)M$), wherein M is conveniently selected from sodium and potassium.

The alkyl sulfate surfactants hereof are water soluble salts or acids of the formula $ROSO_3M$ wherein R preferably is a C_{10} - C_{24} hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C_{10} - C_{18} alkyl component, more preferably a C_{12} - C_{15} alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperidinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like).

The amine

The amines as used herein refer to primary or tertiary amines which is believed to form a mixed micelle with the anionic surfactant and where the carbon chain length of the alkyl group is equal or greater than C₆.

Suitable primary amines for use herein include amines according to the formula R₁R₂R₃N wherein R₁ and R₂ are both H, R₃ is a C₄-C₁₈, preferably C₆-C₁₂ alkyl chain, R₃ alkyl chains may be straight or branched and may be interrupted with up to 12 ethylene oxide moieties, most preferably interrupted with up to 5 ethylene oxide moieties. Preferred amines according to the formula herein above are n-alkyl amines. Suitable amines for use herein may be selected from 1-hexylamine, 1-octylamine, laurylamine, palmitylamine, stearylamine, oleylamine, coconutalkylamine, tallowalkylamine,

Other suitable primary amines include amines according to the formula R₁R₂R₃N wherein R₁ and R₂ are both H ; R₃ is R₄X(CH₂)_n, X is -O-, -C(O)NH- or -NH-, R₄ is a C₄-C₁₈, preferably C₆-C₁₂ alkyl chain and R₄ may be branched or straight, n is between 1 to 5. Preferred amines according to the formula herein above 3-isopropoxypropylamine, 3-(2-methoxyethoxy)-propylamine and 2-(2-aminoethoxy)-ethanol, C₈-C₁₀ octyl oxy propylamine, 2-ethylhexyloxypropylamine, lauryl amido propylamine and coco amido propylamine.

Suitable tertiary amines for use herein include amines tertiary amines having the formula R₁R₂R₃N wherein neither R₁ nor R₂ is H, R₁ and R₂ are C₁-C₈ alkylchains or

$$\text{---}\left(\text{CH}_2\text{---}\overset{\text{R}_5}{\underset{|}{\text{CH}}}\text{---O}\right)_x\text{H}$$
 whereby n is between 2 to 4 and x is between 1 to 6;

R₃ is either a C₄-C₁₈, preferably C₆-C₁₂ alkyl chain, or R₃ is R₄X(CH₂)_n, whereby X is -O-, -C(O)NH- or -NH-, R₄ is a C₄-C₁₈, n is between 1 to 5, and R₅ is H or C₁-C₂ alkyl.

R₃, R₄ are preferably C₆-C₁₂ alkyl chains and may be straight or branched ; R₃ alkyl chains may be interrupted with up to 12 ethylene oxide moieties, most preferably interrupted with up to 5 ethylene oxide moieties.

Suitable tertiary amines for use herein include coconutalkyldimethylamine, dimethyloleylamine, hexadecyltris (ethyleneoxy)dimethylamine, tallowalkylbis(2-hydroxyethyl)amine, stearylbis(2-hydroxyethyl)amine and oleoylbis(2-hydroxyethyl)amine.

Of all of the foregoing amines the preferred materials are the trialkyl amines marketed under the tradename ADOGEN, the long chain alkyldimethyl amines marketed under the tradename ARMEEN and the ethoxylated amines marketed under the tradename ETHOMEEN. The most preferred amines for use in the compositions herein are 1-hexylamine, 1-octylamine, 1-decylamine, 1-dodecylamine. Especially desirable for odor characteristics are n-dodecyldimethylamine (ARMEEN DM12D) and bis(2-hydroxyethyl)coconutalkylamine (ETHOMEEN C/12, BEROL 307) and oleylamine 7 times ethoxylated (BEROL 28), lauryl amido propylamine and coco amido propylamine.

While not intending to be limited by theory, it is believed that the amine cosurfactant and anionic surfactant herein form complexes which enhance packing of the surfactants at the oil/water interface, thereby lowering interfacial tension and improving detergency. The amine surfactants would be at least partially protonated in the product and during the wash thus can form positively charged species capable of complexing with the anionic surfactant.

Another aspect of the invention relates to the surprising finding that low amounts of certain primary and/or tertiary amines of the present invention provide suds control to the detergent compositions formulated therewith. The amounts in which the amines are used for controlling the suds are from 0.1 to 10%, preferably from 0.1 to 5%, most preferably from 0.5 to 4% by weight of the detergent composition.

Detergent ingredients

In another embodiment of the present invention, a liquid detergent composition is provided comprising the surfactant system of the present invention mixed with detergent ingredients. A wide range of surfactants can be used in the detergent composition of the present invention. The detergent compositions according to the present invention will preferably comprise a surfactant system which is substantially free of linear alkylbenzene sulfonate surfactant.

A typical listing of anionic, nonionic, ampholytic and zwitterionic classes, and species of these surfactants, is given in US Patent 3,664,961 issued to Norris on May 23, 1972.

Other suitable anionic surfactants that can be used are alkyl ester sulfonate surfactants including linear esters of C₈-C₂₀ carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO₃ according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula :



wherein R³ is a C₈-C₂₀ hydrocarbyl, preferably an alkyl, or combination thereof, R⁴ is a C₁-C₆ hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R³ is C₁₀-C₁₆ alkyl, and R⁴ is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R³ is C₁₀-C₁₆ alkyl.


Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono-, di- and triethanolamine salts) of soap, C₈-C₂₂ primary or secondary alkanesulfonates, C₈-C₂₄ olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C₈-C₂₄ alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleoyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C₁₂-C₁₈ monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C₆-C₁₂ diesters), sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), and alkyl polyethoxy carboxylates such as those of the formula $RO(CH_2CH_2O)_k-CH_2COO-M^+$ wherein R is a C₈-C₂₂ alkyl, k is an integer from 0 to 10, and M is a soluble salt-forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil. Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 5% to about 25% by weight of such anionic surfactants.

One class of nonionic surfactants useful in the present invention are condensates of ethylene oxide with a hydrophobic moiety to provide a surfactant having an average hydrophilic-lipophilic balance (HLB) in the range from 8 to 17, preferably from 9.5 to 14, more preferably from 12 to 14. The hydrophobic (lipophilic) moiety may be aliphatic or aromatic in nature and the length of the polyoxyethylene group which is condensed with any particular hydrophobic group can be readily adjusted to yield a water-soluble compound having the desired degree of balance between hydrophilic and hydrophobic elements.

Especially preferred nonionic surfactants of this type are the C₉-C₁₅ primary alcohol ethoxylates containing 3-12 moles of ethylene oxide per mole of alcohol, particularly the C₁₂-C₁₅ primary alcohols containing 5-8 moles of ethylene oxide per mole of alcohol.

Another class of nonionic surfactants comprises alkyl polyglucoside compounds of general formula



wherein Z is a moiety derived from glucose; R is a saturated hydrophobic alkyl group that contains from 12 to 18 carbon atoms; t is from 0 to 10 and n is 2 or 3; x is from 1.3 to 4, the compounds including less than 10% unreacted fatty alcohol and less than 50% short chain alkyl polyglucosides. Compounds of this type and their use in detergent are disclosed in EP-B 0 070 077, 0 075 996 and 0 094 118.


Very suitable as nonionic surfactants are poly hydroxy fatty acid amide surfactants of the formula



wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative thereof. Preferably, R¹ is methyl, R² is a straight C₁₁₋₁₅ alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z

is derived from a reducing sugar such as glucose, fructose, maltose, lactose, in a reductive amination reaction.

Highly preferred nonionics are amine oxide surfactants. The compositions of the present invention may comprise amine oxide in accordance with the general formula I:



In general, it can be seen that the structure (I) provides one long-chain moiety $R^1 (EO)_x (PO)_y (BO)_z$ and two short chain moieties, $CH_2 R'$. R' is preferably selected from hydrogen, methyl and $-CH_2OH$. In general R^1 is a primary or branched hydrocarbyl moiety which can be saturated or unsaturated, preferably, R^1 is a primary alkyl moiety. When $x+y+z = 0$, R^1 is a hydrocarbyl moiety having chainlength of from about 8 to about 18. When $x+y+z$ is different from 0, R^1 may be somewhat longer, having a chainlength in the range $C_{12}-C_{24}$. The general formula also encompasses amine oxides wherein $x+y+z = 0$, $R^1 = C_8-C_{18}$, $R' = H$ and $q = 0-2$, preferably 2. These amine oxides are illustrated by C12-14 alkyldimethyl amine oxide, hexadecyl dimethylamine oxide, octadecylamine oxide and their hydrates, especially the dihydrates as disclosed in U.S. Patents 5,075,501 and 5,071,594, incorporated herein by reference.

The invention also encompasses amine oxides wherein $x+y+z$ is different from zero, specifically $x+y+z$ is from about 1 to about 10, R^1 is a primary alkyl group containing 8 to about 24 carbons, preferably from about 12 to about 16 carbon atoms; in these embodiments $y + z$ is preferably 0 and x is preferably from about 1 to about 6, more preferably from about 2 to about 4; EO represents ethyleneoxy; PO represents propyleneoxy; and BO represents butyleneoxy. Such amine oxides can be prepared by conventional synthetic methods, e.g., by the reaction of alkylethoxysulfates with dimethylamine followed by oxidation of the ethoxylated amine with hydrogen peroxide.

Highly preferred amine oxides herein are solids at ambient temperature, more preferably they have melting-points


in the range 30°C to 90°C. Amine oxides suitable for use herein are made commercially by a number of suppliers, including Akzo Chemie, Ethyl Corp., and Procter & Gamble. See McCutcheon's compilation and Kirk-Othmer review article for alternate amine oxide manufacturers. Preferred commercially available amine oxides are the solid, dihydrate ADMOX 16 and ADMOX 18, ADMOX 12 and especially ADMOX 14 from Ethyl Corp.

Preferred embodiments include hexadecyldimethylamine oxide dihydrate, octadecyldimethylamine oxide dihydrate, hexadecyltris(ethyleneoxy)dimethyl-amine oxide, and tetradecyldimethylamine oxide dihydrate.

Whereas in certain of the preferred embodiments $R' = H$, there is some latitude with respect to having R' slightly larger than H . Specifically, the invention further encompasses embodiments wherein $R' = CH_2OH$, such as hexadecylbis(2-hydroxyethyl)amine oxide, ² tallowbis(2-hydroxyethyl)amine oxide, stearylbis(2-hydroxyethyl)amine oxide and oleylbis(2-hydroxyethyl)amine oxide.

When included therein, the laundry detergent compositions of the present invention typically comprise nonionic surfactants in the weight ratio of anionic surfactant to nonionic surfactant from 6:1 to 1:3, preferably from 5:1 to 2:1.

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyldimethylammonium halogenides, and those surfactants having the formula :



wherein R^2 is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-CH_2CH_2-$, $-CH_2CH(CH_3)-$, $-CH_2CH(CH_2OH)-$, $-CH_2CH_2CH_2-$, and mixtures thereof; each R^4 is

selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, benzyl ring structures formed by joining the two R⁴ groups, -CH₂CHOH-CHOHCOR⁶CHOHCH₂OH wherein R⁶ is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R⁵ is the same as R⁴ or is an alkyl chain wherein the total number of carbon atoms of R² plus R⁵ is not more than about 18; each y is from 0 to about 10 and the sum of the y values is from 0 to about 15; and X is any compatible anion.

Preferred cationic surfactants are the water-soluble quaternary ammonium compounds useful in the present composition having the formula :



wherein R₁ is C₈-C₁₆ alkyl, each of R₂, R₃ and R₄ is independently C₁-C₄ alkyl, C₁-C₄ hydroxy alkyl, benzyl, and - (C₂H₄O)_xH where x has a value from 1 to 5, and X is an anion. Not more than one of R₂, R₃ or R₄ should be benzyl.

The preferred alkyl chain length for R₁ is C₁₂-C₁₅ particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis. Preferred groups for R₂R₃ and R₄ are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are :

- coconut trimethyl ammonium chloride or bromide;
- coconut methyl dihydroxyethyl ammonium chloride or bromide;
- decyl triethyl ammonium chloride;
- decyl dimethyl hydroxyethyl ammonium chloride or bromide;
- C₁₂-15 dimethyl hydroxyethyl ammonium chloride or bromide;
- coconut dimethyl hydroxyethyl ammonium chloride or bromide;
- myristyl trimethyl ammonium methyl sulphate;
- lauryl dimethyl benzyl ammonium chloride or bromide;
- lauryl dimethyl (ethenoxy)₄ ammonium chloride or bromide;

choline esters (compounds of formula (i) wherein R_1 is $-\text{CH}_2-\text{O}-\text{C}(\text{O})-\text{C}_{12-14}$ alkyl and $R_2R_3R_4$ are methyl).

Other cationic surfactants useful herein are also described in U.S. Patent 4,228,044, Cambre, issued October 14, 1980.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.5% to about 5%, preferably from about 1% to about 3% by weight of such cationic surfactants.

The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetraacetate, metal ion sequestrants such as aminopolyphosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylenephosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable polycarboxylates builders for use herein include citric acid, preferably in the form of a water-soluble salt, derivatives of succinic acid of the formula $\text{R}-\text{CH}(\text{COOH})\text{CH}_2(\text{COOH})$ wherein R is C_{10-20} alkyl or alkenyl, preferably C_{12-16} , or wherein R can be substituted with hydroxyl, sulfo sulfoxyl or sulfone substituents. Specific examples include lauryl succinate, myristyl succinate, palmityl succinate 2-dodecenylsuccinate, 2-tetradecenyl succinate. Succinate builders are preferably used in the form of their water-soluble salts, including sodium, potassium, ammonium and alkanolammonium salts.

Other suitable polycarboxylates are oxodisuccinates and mixtures of tartrate monosuccinic and tartrate disuccinic acid such as described in US 4,663,071.

Especially for the liquid execution herein, suitable fatty acid builders for use herein are saturated or unsaturated C_{10-18} fatty acids, as well as the corresponding soaps. Preferred saturated species have from 12 to 16 carbon atoms in the alkyl chain. The preferred unsaturated fatty acid is oleic acid.

Other preferred builder system for liquid compositions is based on dodecenyl succinic acid and citric acid.

Detergency builder salts are normally included in amounts of from 3% to 50% by weight of the composition preferably from 5% to 30% and most usually from 5% to 25% by weight.

Other components used in detergent compositions may be employed, such as enzymes and stabilizers therefore, soil-suspending agents, soil-release polymers, abrasives, bactericides, tarnish inhibitors, coloring agents, foam control agents, corrosion inhibitors and perfumes.

Preferably, the liquid compositions according to the present invention are in "concentrated form"; in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. The level of water is less than 50%, preferably less than 30%, more preferably less than 20% of water by weight of the detergent compositions.

Said concentrated products provide advantages to the consumer, who has a product which can be used in lower amounts and to the producer, who has lower shipping costs.

The liquid compositions are especially effective when applied directly to soils and stains in a pretreatment step. The detergent compositions of the present invention can also be used as detergent additive products. Such additive products are intended to supplement or boost the performance of conventional detergent compositions.

The detergent compositions according to the present invention include compositions which are to be used for cleaning of substrates, such as fabrics, fibers, hard surfaces, skin etc., for example hard surface cleaning compositions (with or without abrasives), laundry detergent compositions, automatic and non-automatic dishwashing compositions.

The following examples are meant to exemplify compositions of the present inventions, but are not necessarily meant to limit the scope of the invention.

EXAMPLE I

The following liquid detergent compositions are made :

% by weight of th detergent compositions

	A	B	C	D
C ₁₂ -C ₁₅ Alkyl ethoxylated sulfate	21	11.0	17	16.0
Octylamine	0.5	1	0.5	0.5
C ₁₂ -C ₁₄ alkyldimethyl amine oxide	3	-	2	2
C ₁₂ -C ₁₅ Alkyl sulfate	5	17.0	-	10
C ₁₂ -C ₁₄ N-methyl glucamide	5	5.0	4.0	5
C ₁₂ -C ₁₄ fatty alcohol ethoxylate	1	1	1	1
C ₁₂ -C ₁₆ Fatty acid	7	7	4	6
Citric acid anhydrous	4.5	4.5	3	4
Diethylene triamine penta methylene phosphonic acid	1.0	1.0	2.0	2.0
Monoethanolamine	8	8	0.5	9
Sodiumhydroxide	1	1	2.5	1
Propanediol	12.7	14.5	13.1	10.0
Ethanol	1.8	1.8	4.7	5.4
Amylase (300KNU/g)	0.1	0.1	0.1	0.1
Lipolase(100KNU/g)	0.15	0.15	0.15	0.15
Protease (34g/l)	0.5	0.5	0.5	0.5
Endo-A (5000 CEVU/g)	0.05	0.05	0.05	0.05
Carezyme (5000 CEVU/g)	0.09	0.09	0.09	0.09
Terephthalate-based polymer	0.5	0.5	-	0.3
Boric acid	2.4	2.4	2.8	2.8
Sodium xylene sulfonate	-	-	3	-
DC 3225C	0.03	0.04	0.03	0.04
2-butyl-octanol	0.03	0.04	0.04	0.03
Branched silicone	0.3	0.3	0.3	0.3
Water & Minors	-----up to 100%-----			


The above liquid detergent compositions (A-D) are found to be very efficient in the removal of greasy/oily soils under various usage conditions while having a controlled suds profile. The same results were obtained when octylamine was replaced in Compositions (A-D) by Bis(2-hydroxyethyl)coco amine, Oleyl amine 7EO, lauryl amido propylamine and coco amido propylamine.

What is claimed is:

1. A liquid detergent compositions comprising anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates, characterized in that said detergent composition further comprises an amine selected from

i) primary amines having the formula $R_1R_2R_3N$ wherein R_1 and R_2 are both H ; and wherein R_3 is C_4-C_{18} or $R_4X(CH_2)_n$ whereby X is -O-, -C(O)NH- or -NH-, R_4 is a C_4-C_{18} , n is between 1 to 5;

ii) tertiary amines having the formula $R_1R_2R_3N$ wherein neither R_1 nor R_2 is H, R_1 and R_2 are C_1-C_8 alkylchains or



whereby n is between 1 to 5 and x is between 1 to 5, R_3 is either C_4-C_{18} or $R_4X(CH_2)_n$ whereby X is -O-, -C(O)NH- or -NH-, R_4 is a C_6-C_{18} , n is between 1 to 5, and R_5 is H or C_1-C_2 alkyl; and;

iii) mixtures of said primary and tertiary amines.


2. A liquid detergent composition according to claim 2 wherein said primary amine is selected from n-alkylamines, C_8-C_{14} alkyl amidopropyl amines, $C_{10}-C_{12}$ alkyl propylene diamines, and C_8-C_{12} alkyl oxypropyl amines..

3. A liquid detergent composition according to claim 2 wherein said n-alkylamine is 1-octylamine.

4. A liquid detergent composition according to claim 1 wherein the amine is selected from the tertiary amine wherein R_1 , R_2

are both $-\left(\text{CH}_2-\overset{\text{R}_5}{\underset{|}{\text{CH}}}-\text{O}\right)_x\text{H}$ whereby n is between 1 to 5 and x is between 1 to 5 and wherein R_3 is C_4-C_{18} alkyl chain and wherein R_5 is H or CH_3 .

5. A liquid detergent composition according to Claim 1 wherein the amine is a tertiary amine selected from C_4-C_{12} alkyl dimethyl amines and amines of the formula



wherein R_4 is C_8 - C_{12} alkyl and n is 2 or 3.

6. A liquid detergent composition according to claims 1-5 wherein the weight ratio of the primary or/and tertiary amine to the alkyl alkoxyated sulfates and alkyl sulfates is from 1:1 to 1:250, more preferably from 1:5 to 1:100 and most preferably from 1:10 to 1:40.

7. A liquid detergent composition according to Claim 6 wherein the anionic surfactants are selected from the alkyl ethoxylated sulfate and C_{12} - C_{15} alkyl sulfate.

8. A liquid detergent composition according to Claims 1-7 further comprising a nonionic surfactant selected from the polyhydroxy fatty acid amides and/or amine oxides.

9. A liquid detergent composition according to claim 8 wherein said amine oxide is C_{12} - C_{14} alkyl dimethyl amine oxide.

10. A liquid detergent composition according to Claims 1-9 further comprising other surfactants, builders, enzymes and other conventional detergent ingredients.

11. A liquid detergent composition according to Claims 1-10 which is free of linear alkylbenzene sulfonate surfactant.

12. Use of the primary amines and/or tertiary amines as defined in claims 1-5 in detergent compositions for the controlling of suds

13. Use of a liquid detergent composition according to Claims 1-11 for pretreatment of fabrics.

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 94/11779A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C11D1/65 C11D3/30 C11D1/86

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP,A,0 203 660 (UNILEVER NV ;UNILEVER PLC (GB)) 3 December 1986 see page 2, column 2, line 14 - page 4, column 5, line 44; claims; tables 1,2 ---	1,2,5-7, 10,13 8
X	FR,A,2 329 746 (PROCTER & GAMBLE EUROP) 27 May 1977 see page 22, line 26 - line 37; examples 4,5,9,13,15 ---	1,2,4-7, 10
A	GB,A,1 375 639 (PROCTER & GAMBLE) 27 November 1974 see claims ---	1,2,4-7, 10
A	US,A,3 853 779 (INAMORATO J ET AL) 10 December 1974 see column 1, line 25 - line 36; claims -----	1,12

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

- * "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- * "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- * "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * "&" document member of the same patent family

Date of the actual completion of the international search

16 June 1995

Date of mailing of the international search report

21.06.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Grittern, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/11779

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0203660	03-12-86	AU-B- 572024	28-04-88
		AU-A- 5797386	04-12-86
		JP-B- 6037635	18-05-94
		JP-A- 61283698	13-12-86

FR-A-2329746	27-05-77	BE-A- 847843	29-04-77
		DE-A- 2648304	05-05-77
		JP-A- 52086410	18-07-77
		NL-A- 7612025	03-05-77

GB-A-1375639	27-11-74	NONE	

US-A-3853779	10-12-74	AU-A- 5503373	07-11-74
		CA-A- 992833	13-07-76
		DE-A- 2323326	03-01-74
		FR-A,B 2187898	18-01-74

THIS PAGE BLANK (USPTO)